

Properties of Particles Obeying Ambiguous Statistics

M.V. Medvedev^{*,†}

Physics Department, University of California at San Diego, La Jolla, California 92093.

A new class of identical particles which may exhibit both Bose and Fermi statistics with respective probabilities p_b and p_f is introduced. Such an uncertainty may be either an intrinsic property of a particle or can be viewed as an “experimental uncertainty”. Statistical equivalence of such particles and particles obeying parastatistics of infinite order is shown. Generalized statistical distributions are derived and statistical and thermodynamical properties of an ideal gas of the particles are investigated. The physical nature of such particles and the implications of this investigation for the statistics of extremal black holes are discussed.

05.30.-d, 04.70.Dy, 14.80.Pb, 05.70.Ce

Historically, the first attempt to generalize quantum Bose and Fermi statistics was made by Gentile [1], who proposed statistics in which up to k particles are allowed to occupy a single quantum state (instead of one for Fermi and infinitely many for Bose cases). Further generalizations were mostly on deformations of commutation relations for particle creation and annihilation operators [2–5]. A special case (the so called “infinite” statistics) is rather interesting in that the particles which obey this statistics have nonnegative square norms [6] for $|q|^2 \leq 1$, while the observables have nonlocal properties. The commutation relations are simply [4,5]:

$$a_i a_j^\dagger - q a_j^\dagger a_i = \delta_{ij}, \quad (1)$$

where q is either a real or complex parameter, $a_j^\dagger$ and a_j are the particle creation and annihilation operators, and δ_{ij} is the Kronecker’s delta. Stimulated by recent experimental observation of the *fractional quantum Hall effect* [7] in a two-dimensional electron gas, various nontraditional statistics have been proposed [8–11]. Most of them are based on generalizations of particle wave-function permutation symmetries. The concept of anyons, which are charge carriers in the fractional quantum Hall effect, is essentially two-dimensional [8]. However, in the strong magnetic field with fluxes antiparallel to the ambient field and in the lowest Landau level, anyons are shown [12] to obey the one-dimensional (though valid in any dimensions) Haldane’s exclusion principle [11]. At last, the equivalence of the anyon statistics (in the above case) and q -on statistics, Eq. (1), with $q = e^{i\alpha\pi}$ (α is the measure of Haldane’s/Pauli blocking and $\alpha\pi$ is exactly the wave-function phase-shift due to permutation of any two particles) was proven [13] via the properties of the N -anyon permutation group.

Statistical and thermodynamical properties of systems of particles obeying fractional statistics (i.e., complex q such as $|q|^2 = 1$) were extensively studied in the last several years [10,12,14], whereas these properties of particles with real q , $-1 < q < 1$, were not investigated (with some special exceptions [15,16]). One should mention a special case of the infinite statistics [4] with $q = 0$ which

was suggested [3] to be equivalent to the statistics of non-identical particles (quantum Boltzmann statistics). One should note here that it was also recently suggested [17] that extremal black holes should obey such a statistics. This issue will be discussed below.

In this work we investigate a system of particles obeying ambiguous statistics which are (statistically) equivalent to those obeying the q -deformed statistics of real q . The partition function of a free gas of such particles, however, differs from that of a free q -on gas. The last is shown [16] to be independent of the deformation parameter q . In fact, the second quantized approach used in [16] does not take into account the dynamics of the internal degrees of freedom. Thus, the q -ons are always distinguishable particles. Our model, however, is (probably) identical to the case when the internal degrees of freedom are in the thermodynamic equilibrium with the external degrees of freedom (at least for $q = 0$), and thus the particles are *not completely distinguishable*. (Note, the particles are indistinguishable provided they are in the same internal state, even if they have infinite number of internal degrees of freedom.) Consequently, the quantum Boltzmann distribution ($q = 0$) is different from the classical Boltzmann.

Unlike the theories mentioned above, we admit only “primary” Bose-Einstein and Fermi-Dirac statistics as existing. Assume now that a particle is neither a pure boson or pure fermion. Let another particle, which interacts with the first one, plays a role of an external observer. During the interaction, it performs a measurement at the first particle and identifies it as either a boson or a fermion with respective probabilities p_b and p_f . According to the result of this measurement, it interacts with the first particle as if the last is a boson or fermion, respectively. The first particle, of course, is the observer for the second particle, thus the process is symmetric. Note, that $p_b + p_f$ is not necessarily equal to one, and if not, it means that the second particle (observer) does not detect the first particle. The probability of this is $1 - p_b - p_f$. The “statistical uncertainty” introduced above may be either the intrinsic property of a particle itself or the “experi-

mental uncertainty” of the measurement process. This concept of “imperfect measurement” is similar to that proposed in order to resolve the causality paradox in superluminal particle (tachyon) systems [18]. This model can be interpreted in another manner. Assume a particle can oscillate between two types of statistics, then the model we propose represents a system of such particles averaged over time-scales much larger than the oscillation period. The probabilities p_b and p_f , thus, are those portions of time during which a particle resides in a Fermi- or Bose-type state. The model of this kind is relevant to systems of elementary particles which have mutual supersymmetrical partners (e.g., a photon and photino, etc.) when transitions between these supersymmetrical states can occur [19]. With some modifications, this model can also be applied to systems of particles with changing flavor, e.g., quarks, gluons, neutrinos, etc.

There is a conjectured relation between the particles of the ambiguous statistical type and the q -deformed infinite statistics. Let b_j and $b_j^\dagger$ are the annihilation and creation operators for such a particle. The particle exhibit bosonic properties with the probability p_b and fermionic ones with p_f . Thus, a bilinear commutation relation for b_j and $b_j^\dagger$ is of boson type with the probability p_b^2 , of fermion type with the probability p_f^2 , and the two particles are nonidentical with the probability $2p_b p_f$, that is interchange of their positions result in another wavefunction, i.e., $b_i b_j^\dagger = \delta_{ij}$. We write

$$(p_b^2 + p_f^2 + 2p_b p_f) b_i b_j^\dagger - (p_b^2 - p_f^2) b_j^\dagger b_i = (p_b^2 + p_f^2 + 2p_b p_f) \delta_{ij}. \quad (2)$$

This equation coincides with Eq. (1), and the deformation parameter is $q = (p_b - p_f)/(p_b + p_f)$. In the rest of this letter we derive statistical distributions for identical particles which have ambiguous quantum exclusion properties and investigate some thermodynamical properties of an ideal gas of such particles.

A grand partition function for a system of particles with properties defined by a stochastic label with a known probability distribution is a sum over all possible realizations (with a weight factor) of the partition functions corresponding to each realization. In each realization, the system effectively consists of k bosons and $N - k$ fermions (N is the total number of particles), while the probability of this realization is $p_b^k p_f^{N-k}$. The total number of states of the system is

$$W = \prod_j \sum_{k=0}^{N_j} \binom{N_j}{k} w_{b_j}(k) w_{f_j}(N_j - k) p_b^k p_f^{N_j - k}, \quad (3)$$

where the weight factor $\binom{N}{k} = \frac{N!}{k!(N-k)!}$ represents the number of ways to arrange N identical particles in two groups of respectively k and $N - k$ particles each. Here $w_b(m)$ and $w_f(m)$ are the numbers of quantum states

of m identical particles occupying a group of G states, respectively, for bosons and fermions,

$$w_b(m) = \frac{(G + m - 1)!}{m!(G - 1)!} \quad \text{and} \quad w_f(m) = \frac{G!}{m!(G - m)!}. \quad (4)$$

The partial partition function (expression under the product sign) can be simplified and rewritten in terms of a generalized hypergeometric function

$$W_j = p_f^{N_j} \frac{G_j}{N_j} \sum_{k=0}^{N_j} \binom{N_j}{k}^2 \binom{G_j - 1 + k}{N_j - 1} \left(\frac{p_b}{p_f}\right)^k \quad (5a)$$

$$= p_f^{N_j} G_j \frac{G_j!}{N_j!(G_j - N_j + 1)!} \cdot {}_3F_2 \left(-N_j, -N_j, G_j; 1, G_j - N_j + 1; \frac{p_b}{p_f} \right). \quad (5b)$$

To proceed further, we assume the number of particles in a system to be very large, and use Stirling approximation for the factorials and Laplace’s approximation method to evaluate the sum. It is enough to keep the leading term, only. Then one can write Eq. (5) as follows

$$W_j \simeq \binom{N_j}{k_{0j}}^2 \binom{G_j + k_{0j}}{N_j} p_b^{k_{0j}} p_f^{N_j - k_{0j}}, \quad (6)$$

where k_{0j} is the solution of the equation

$$\left(\frac{N_j - k_{0j}}{k_{0j}} \right)^2 \left(\frac{G_j + k_{0j}}{G_j - N_j + k_{0j}} \right) \frac{p_b}{p_f} = 1. \quad (7)$$

Thus one can infer that the partition function, although defined by all realizations, has its major contribution from that realization in which exactly k_0 bosons and $N - k_0$ fermions effectively exist, where k_0 is defined by the probability ratio p_b/p_f alone. (Note, it is not a mixture of species, there is only one physically existing species.) The entropy of the system is, as usual, $S = \ln W$:

$$S \simeq \sum_j G_j \left\{ x_j \ln(p_b/p_f) + n_j \ln p_f + (x_j + 1) \ln(x_j + 1) + n_j \ln n_j - (1 - n_j + x_j) \ln(1 - n_j + x_j) - 2[x_j \ln x_j + (n_j - x_j) \ln(n_j - x_j)] \right\}, \quad (8)$$

where $n_j = N_j/G_j$ is the occupation number and $x_j = k_{0j}/G_j$. Other thermodynamic functions follow straightforwardly. The most probable distribution subject to the constraints that the total number of particles and the total energy of the system be conserved, is determined by the condition

$$\frac{\partial}{\partial N_j} \left[S - \alpha \sum_j N_j - \beta \sum_j \epsilon_j N_j \right] = 0. \quad (9)$$

From this equation and Eqs. (7), (8), we arrive at

$$\frac{n_j(1 - n_j + x_j)}{(n_j - x_j)^2} p_f = \rho, \quad (10)$$

where $\rho = \exp\{(\epsilon_j - \mu)/T\}$, $\mu \equiv -\alpha T$ is the chemical potential, and $T \equiv 1/\beta$ is the temperature of the gas. Equations (7) and (10) define the occupation number $n(\epsilon_j)$ as a function of energy. Upon simple mathematical manipulations, we obtain

$$x_j = \frac{n_j(\rho + \delta) - \delta}{2\rho + \delta} \quad (11)$$

and

$$(\rho + \delta)(\rho + p_f)(\rho - p_b)n_j^2 - 2\rho(\rho + \delta)\sigma n_j + \rho\delta^2 = 0, \quad (12)$$

where $\delta = p_f - p_b$ and $\sigma = p_f + p_b$. Note, the fraction x_j/n_j is the number of bosons in the system with respect to the total number of particles, i.e., $0 \leq x_j/n_j \leq 1$. The last equation can be explicitly solved to obtain

$$n_j = \frac{\sigma\rho}{(\rho + p_f)(\rho - p_b)} \left[1 + \sqrt{1 - \frac{\delta^2}{\sigma^2} \frac{(\rho + p_f)(\rho - p_b)}{\rho(\rho + \delta)}} \right]. \quad (13)$$

Here we omit the negative sign of the root, because it does not satisfy [together with Eq. (11)] the condition $0 \leq x_j/n_j \leq 1$. In order to n_j be positive number for all values of ϵ_j , the chemical potential should not exceed its maximum value $\mu_{max} = -T \ln p_b$. Eq. (13) is a generalized energy distribution for the class of particles obeying the ambiguous statistics as a function of two parameters p_b and p_f which define the deviation from “primary” Bose or Fermi statistics. The generalized distribution recovers Bose-Einstein and Fermi-Dirac ones in the limiting cases

$p_b = 1, p_f = 0$ and $p_b = 0, p_f = 1$, respectively. The case $p_b = p_f = \sigma/2$ is that of the quantum Boltzmann statistics since the deformation parameter $q = 0$. In this case

$$n_j = 2\sigma \left(\rho - \frac{\sigma^2}{4} \frac{1}{\rho} \right)^{-1}, \quad (14)$$

which is a quantum analog of the Boltzmann distribution. It is remarkable that although particles can, in principle, be distinguished by their internal states (they have an infinite number of internal degrees of freedom), the occupation number of these states is also energy and temperature dependent. Thus, the particles, in fact, are *not completely* distinguishable. Hence, some trend towards the familiar statistical (and thermodynamical) properties of identical particles may be discerned. It is also interesting that Eq. (14) reduces to the classical Boltzmann distribution $\sim \rho^{-1}$ for a weakly interacting gas, $\sigma \rightarrow 0$ (remember, particles do not “feel” each other with the probability $1 - p_b - p_f = 1 - \sigma$). The factor 2σ is not significant since it just renormalizes μ .

All thermodynamic properties can be derived straightforwardly [20]. The chemical potential μ is defined implicitly by the equation for the total particle number

$$N = \frac{Vm^{2/3}}{\sqrt{2}\pi^2\hbar^3} \int_0^\infty \sqrt{\epsilon} n(\epsilon) d\epsilon, \quad (15)$$

where V is the volume of the gas, m is the particle mass, $\hbar$ is the Plank’s constant. Then the equation of state can be found in a parametric form from the thermodynamic potential $\Omega = -PV$,

$$\Omega = -\frac{2}{3} \frac{Vm^{2/3}}{\sqrt{2}\pi^2\hbar^3} \int_0^\infty \epsilon^{3/2} n(\epsilon) d\epsilon, \quad (16)$$

In the classical Boltzmann limit $\exp\{\mu/T\} \ll 1$ the equation of state [20] expanded to second order is

$$PV = NT \left\{ 1 + \frac{\delta}{(\sigma + 2\sqrt{p_b p_f})} \frac{\pi^{3/2}}{2} \frac{N\hbar^3}{V(mT)^{3/2}} - \left[\frac{(\delta^2 + p_b p_f)}{(\sigma + 2\sqrt{p_b p_f})^2} + \frac{\delta^2 \sqrt{p_b p_f}}{4(\sigma + 2\sqrt{p_b p_f})^3} \right] \frac{16\pi^3}{3^{5/2}} \frac{N^2 \hbar^6}{V^2(mT)^3} \right\}. \quad (17)$$

There is an effective attraction between particles of the gas when $p_b > p_f$ and effective repulsion when $p_b < p_f$ (the second term overcomes the last one), as expected. In the case of the quantum Boltzmann statistics, $\delta = 0$, the second term vanishes, but a residual attraction still exists (the last quadratic term).

As was mentioned above, extremal black holes are shown to obey the infinite statistics with the deformation parameter $q = 0$ [17]. Let’s consider a (“gedanken”) gas of charged, nonrotating, extremal black holes, and assume their charges are of the same sign. Then, by definition, $M^2 = Q^2$ (with M and Q being the mass and

charge of a black hole). In the limit of large black hole separation (i.e., low-density, ideal gas approximation, no relativistic effects), gravitational attraction of “particles” completely cancels electrostatic repulsion. Thus, the system at hand is a gas of noninteracting particles (neutrals). The thermodynamical properties of such a gas are governed by the particle statistics properties, alone. According to Eq. (17) (with $p_b = p_f = \sigma/2, \delta = 0$), the nonrotating, extremal black holes will experience *weak statistical attraction*. Hence, one may expect that black holes will form clusters, for instance, the ultimate size of which is controlled by the gas temperature (i.e., the

velocity dispersion of black holes) and completely independent of *any* black hole parameters (e.g., their masses and charges).

To summarize, we introduced a new class of particles which may fluctuate between Bose- and Fermi-type statistics or may have uncertainty in their statistics. A generalized distribution function is derived. For a particular case of the quantum Boltzmann statistics, the distribution function (i.e., occupation numbers) is different from the classical Boltzmann case. An equation of state in the classical Boltzmann limit is derived. The relation of this type of statistics to the existing generalizations of Bose and Fermi statistics has been ascertained. The implication of these results obtained to the ensemble of extremal black holes is discussed.

I am grateful to P.H. Diamond for useful discussions. This work was supported by DoE grant No. DEFG0388ER53275.

-
- * E-mail address: mmedvedev@ucsd.edu,
Internet:
<http://sdphpd.ucsd.edu/~medvedev/mm.html>
- † Also: Russian Research Center “Kurchatov Institute”,
Institute for Nuclear Fusion, Moscow 123182, RUSSIA.
- [1] G. Gentile, *Nuovo Cimento* **17**, 493 (1940);
[2] H.S. Green, *Phys. Rev.* **90**, 270 (1952); O.W. Greenberg,
Phys. Rev. Lett. **13**, 598 (1964); O.W. Greenberg, in
Mathematical Theory of Elementary Particles, Proceedings of the conference, MIT Press (1966).

- [3] A.B. Govorkov, *Phys. Part. Nucl.* **24**, 565 (1993).
[4] O.W. Greenberg, *Phys. Rev. Lett.* **64**, 705 (1990).
[5] O.W. Greenberg, *Phys. Rev. D* **43**, 4111 (1991).
[6] R.N. Mohapatra, *Phys. Lett. B* **242**, 407 (1990).
[7] D.C. Tsui, H.L. Stromer, and A.C. Gossard, *Phys. Rev. Lett.* **48**, 1559 (1982); H.L. Stromer, A. Chang, D.C. Tsui, J.C.M. Hwang, A.C. Gossard, and W. Wiegmann, *Phys. Rev. Lett.* **50**, 1953 (1983).
[8] J.M. Leinaas and J. Myrheim, *Nuovo Cimento B* **37**, 1 (1977); F. Wilczek, *Phys. Rev. Lett.* **49**, 957 (1982).
[9] B.I. Halperin, *Phys. Rev. Lett.* **52**, 1583 (1983); R.B. Laughlin, *Phys. Rev. Lett.* **60**, 2677 (1988); F. Calogero, *J. Math. Phys. (N.Y.)* **10**, 2191 (1969); **10** 2197 (1969); B. Sutherland, *J. Math. Phys. (N.Y.)* **12**, 246 (1971); **12** 251 (1971); *Phys. Rev. Lett.* **34**, 1083 (1975).
[10] M.V.N. Murthy and R.Shankar, *Phys. Rev. Lett.* **73**, 3331 (1994).
[11] F.D.M. Haldane, *Phys. Rev. Lett.* **67**, 937 (1994).
[12] A. Dasnieres de Veigy and S. Ouvry, *Phys. Rev. Lett.* **72**, 600 (1994); Y. Wu, *Phys. Rev. Lett.* **73**, 922 (1994).
[13] G.A. Goldin and D.H. Sharp, *Phys. Rev. Lett.* **76**, 1183 (1996).
[14] S.B. Isakov, *Phys. Rev. Lett.* **73**, 2150 (1994); D. Sen and R.K. Bhaduri, *Phys. Rev. Lett.* **74**, 3912 (1994); A. Smerzi, *Phys. Rev. Lett.* **76**, 559 (1995).
[15] A. Inomata and S. Kirchner, *Phys. Lett. A* **222**, 213 (1996); S. Dalton and A. Inomata, *Phys. Lett. A* **199**, 315 (1995).
[16] R.F. Werner, *Phys. Rev. D* **48**, 2929 (1993).
[17] A. Strominger, *Phys. Rev. Lett.* **71**, 3397 (1993).
[18] L.S. Schulman, *Am. J. Phys.* **30**, 481 (1970).
[19] G. Kane, *Modern Elementary Particle Physics*, (Addison-Wesley Pub. Company, Inc., 1987).
[20] L.D. Landau and E.M. Lifshitz, *Statistical Physics*, (Pergamon, Oxford, 1980).